



**Bharati Vidyapeeth University Medical College
Hospital & Research Centre, Pune
Antimicrobial Policy
and
Antimicrobial Stewardship Program
2024-2025
Version - 10.0**



Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology 	Dr M S Modak Prof Microbiology 	Dr AK Verma Quality Department of Med Dir 	Dr SK Dalwadi Medical Director 



SR NO	CONTENTS	PAGE NO
Prelude	List of amendments in Version 10	3
1	Introduction	4
2.	Clinical pathway	5
3.	Antimicrobial Stewardship Goals Workflow, Metrics	6-7
4.	List of restricted antimicrobials and Types of antimicrobial resistant organisms	8-9
5.	Common isolates from various samples and their antibiogram for the year: 2022 Critical care Medicine Surgery Neonatology Obst Gynae Orthopaedics Paediatrics& PICU Oncology OPD (All) Index of multidrug resistance	10-13 14-17 18- 20 21-23 24-26 27-29 30- 32 33-36 37-39 40
6.	Antimicrobial therapy in hospitalized patients	41
7.	Empiric Antimicrobial choice in various clinical conditions according to patient category, Definitive therapy	42-65
8.	Surgical Antimicrobial prophylaxis/Gastrointestinal endoscopy prophylaxis	63-65
9.	Febrile Neutropenia Cardiac device implant prophylaxis	69-71 72
10.	Appendix 1 : Commonly used antimicrobials , their spectrum of activity and side effects	73-76
11.	Appendix 2 Duration of therapy of common conditions	77
12.	Appendix 3 : Antimicrobial Agent form	78
13.	Appendix 4 : Sample collection methods	79-81
14.	References	82

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



List of amendments in Version 10.0

1. Isolate listing and Antibiogram for the year 2023
2. Antibiogram : Sample type Pus has been separated into Wound and Tissue for Surgical and Orthopaedic specialities
3. Added : OPD cases of Acute gastroenteritis suggested antibiotic if required
4. UTI in Obst-Gynae : Added Table 4a Cefixime and Fosfomycin after 20wk pregnancy (added reference No 9)
5. Updated vancomycin dosing for surgical prophylaxis in Table 16
6. Added Table 20 for Cardiac device implantation prophylaxis (added reference No 10)
7. Renumbered older table No 20 to Table 21 (Spectrum of commonly used antimicrobials)
8. Added sample collection method for High Vaginal Swab Appx 4

Bharati Hospital and Research Centre

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



1. Introduction -

Over the last 60 years antimicrobials have been widely used to treat infectious diseases. Their indiscriminate use has led to resistance developing to almost all known antimicrobials. Antimicrobial resistance has become widespread not only in hospitals but also in the community.

A rational antimicrobial policy and antimicrobial stewardship is a must for all hospitals and is mandated by the Ministry of Health and Family Welfare through its document “National Policy for Containment of Antimicrobial Resistance, India”. The purpose of this document is to provide a guide for rational antimicrobial use at Bharati Hospital based on local patterns of antimicrobial susceptibility.

DEFINITIONS

Antimicrobial agent (Antibiotic): Any agent, which has a potential for or is used with an intention of affecting microbial growth inside or on the human body. This includes antibacterial, antifungal, antiviral and anti-parasite agents.

Antimicrobial Stewardship: A set of coordinated activities that includes appropriate selection of antimicrobial agent, dosing, route and duration of antimicrobial therapy. The primary goal of antimicrobial stewardship is to optimize clinical outcomes while minimizing unintended consequences of antimicrobial use, including toxicity, the selection of pathogenic organisms and the emergence of resistance.

Surgical Antimicrobial Prophylaxis/Prophylactic anti-microbial agents: Administration of an antibiotic or antimicrobial agent prior to the commencement of a surgical procedure and appropriate re-administration of the agent during prolonged surgery.

Empiric Antibiotic/Antimicrobial therapy: This is an early institution of antimicrobial therapy pending the results of culture and / or other relevant investigation and clinical response, in patients who have an illness and in whom there is an expectation of an infectious cause, the treatment being directed against the most likely microbial agent(s) in that particular episode.

Organism directed Antimicrobial Therapy: Usage of antimicrobial agent against infection by specific microorganisms which have been confirmed by culture of appropriate samples.

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



2. Clinical Pathway

1. Resident of respective department will assess patient for symptoms and signs of infection, including laboratory evidence of infection.
2. He/she will document appropriately on the culture requisition form.
 - suspected cause/site of infection,
 - possibly community (CA)/hospital acquired(HA)
 - patient type (types 1-3 described below)
3. Appropriate site cultures and blood cultures will be sent according to HICC protocol.
4. Antimicrobial will be chosen according to antimicrobial guide after informing lecturer on call and checking for allergy risks.
5. Any deviation from the policy will be documented along with the reason for deviation.
6. Some antimicrobials will be part of the restricted formulary and use of these “ALERT” antimicrobials will requires infectious disease/ critical care (ICU/PICU/NICU) consult. These include;
Piperacillin/Tazobactam, Carbapenems, Ceftazidime avibactam, Aztreonam, Polymixins, Linezolid, Teicoplanin, Vancomycin, Echinocandins, Voriconazole, Amphotericin B
7. Clinical response will be followed.
8. Once culture reports are available (Day 2 – Day 4) antimicrobial is to be de-escalated (if possible) and duration of therapy is to be specified if not already done so.
9. Antimicrobial prescription should have a record of the day and expected duration of antimicrobials in the left-hand margin of the drug chart, eg D4/7
10. Infection control team will fill antimicrobial audit form and conduct regular department wise audits.
11. Findings of the audit will drive improvement in antimicrobial use.

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



3. Antimicrobial Stewardship



- A set of coordinated strategies to improve the use of **antimicrobials**

Goal


- Enhancing patient health outcomes
- Reducing resistance to **antibiotics**
- Decreasing unnecessary costs

Bharati Hospital and R

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	---	--	--



Workflow of Antimicrobial stewardship



Metrics used in AMSP

- Days of therapy
- **Cost metrics**
- Average length of stay
- **Mortality rate**
- Acceptance of intervention
- **Resistance pattern**
- Comparison of HAI with ABX consumption rate

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



4. List of Restricted Antimicrobials

These will not be prescribed without obtaining concurrence of HoD/HoU ;

1. Piperacillin tazobactam
2. Carbapenems
3. Linezolid
4. Vancomycin
5. Teicoplanin
6. Daptomycin
7. Tigecycline
8. Polypeptides : Polymixin, Colistin
9. Echinocandins : Caspofungin, Micafungin
10. Ceftazidime avibactam
11. Aztreonam

Common antimicrobial resistant organisms :

Extended spectrum beta-lactamase producers (ESBL)

These are Gram negative organisms (GNB) like E coli & Klebsiella, which are resistant to the penicillins; first-, second and third-generation cephalosporins; In addition, the plasmids bearing genes encoding ESBLs frequently also carry genes encoding resistance to other antimicrobial agents, such as aminoglycosides, trimethoprim, sulphonamides, tetracyclines and chloramphenicol. They remain susceptible to beta lactam- beta lactamase inhibitor combinations and carbapenems.

Amp C beta lactamases

These are inducible beta lactamases produced by certain organisms after exposure to cephalosporins. The organisms are resistant to the penicillins; first-, second- and third-generation cephalosporins and beta-lactam-beta lactamase inhibitor combinations. They may remain susceptible to cefepime and carbapenems. Seen in *Serratia*, *Pseudomonas*, *Proteus*, *Citrobacter* and *Enterobacter* spp.

Carbapenemase producers:

These are Gram negative organisms resistant to the Carbapenems and almost all beta-lactam antimicrobials except monobactams. Colistin and polymyxins are currently used for these organisms.

Methicillin Resistant Staphylococcus aureus (MRSA)

These are resistant to all beta lactam antimicrobials (Penicillins, BL-BLI, Cephalosporins, monobactams and Carbapenems.)

Vancomycin Resistant Enterococcus (VRE):

These isolates are resistant to Vancomycin, Teicoplanin but susceptible to linezolid.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



MDR (Multi-drug resistance):

Isolates resistant to representatives of three or more classes of antimicrobial agents,

XDR (Extensive drug resistance):

Isolates resistant to all but one or two classes

PDR (Pan drug resistance):

Isolates resistant to all classes of antimicrobial agents available

Bharati Hospital and Research Centre






Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



5. Organisms commonly isolated and antibiogram: Area wise

INTENSIVE CARE UNIT : TOP 10 ISOLATES BY SAMPLE

SPECIMEN TYPE	OVERALL NO. OF ISOLATES
BLOOD	442
PUS	104
RESPIRATORY	108
URINE	137
CSF	12



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Dilwahi Medical Director
--	---	--	--



INTENSIVE CARE UNIT ANTIBIOGRAM GRAM NEGATIVE

Percentage susceptibility gram negative organism 2023

Location	Specimen type	Organism	Number of patients	LEVOFLOXACIN	CIPROFLOXACIN	COTRIAMOXAZOLE	AMOX/CLAV	Cefuroxime	CEFTAZIDIME	CEFTRIAXONE	CEFOTAXIME	CEFEPIME	CEF/SUL	PIPTAZ	AMIKACIN	GENTAMICIN	ERTAPENEM	MEROPENEM	IMIPENEM	TIGECYCLINE	FOSFOMYCIN	MINOCYCLINE	COLISTIN
SSICU	Blood	Klebsiella pneumoniae	56	0	27	44	29	36		27		34	36	36	47	47	44	46	47		80		100
		Escherichia coli	49		6	50	48	63		25		42	63	63	85	77	73	85	85		98		100
		Acinetobacter baumannii	30		10	33		10		10		7	10	10	10	13		13	10				100
		Burkholderia cepacia	12	80	0	90		20	60	0	0	0	20	20	40	20		80	20				100
	Urine	Escherichia coli	50		4	58	38	58		24		30	58	58	80	74	74	78	76		98		100
		Klebsiella pneumoniae	29		14	55	28	28		21		28	28	28	38	45	41	45	41		79		100
	Respiratory	Acinetobacter baumannii	38		5	29		5		3		5	5	5	5	8		5	5				100
		Klebsiella pneumoniae	17		31	56	38	31		6		19	31	31	38	38	50	44	44		81		100
	Pus	Escherichia coli	21		0	43	19	0		5	0	24	57	57	91	91	62	71	71		91		100
		Klebsiella pneumoniae	14		14	50	29	14		14	14	21	36	36	43	43	36	36	36	36	43		100
		Acinetobacter baumannii	13		0	8				0	0	0	0	0	0	0		0	0		0		92
	Other	Acinetobacter baumannii	10		10	30		10		10		10	10	10	10	10		10	10				100
		Klebsiella pneumoniae	7		14	71	29	43		29		57	43	43	71	71	71	71	57	57	100		100
		Escherichia coli	5		0	50	50	60		0		20	60	60	50	0	50	50	80		100		100
	Fluid	Escherichia coli	12		10	40	36	67		20		58	67	67	90	80	80	90	92		91		100
	Cerebrospinal fluid	Pseudomonas aeruginosa	2	50				100				100	100	100				50	0				100
		Cronobacter sakazakii	1		0	100	100			###						100	100	100	100		100		

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Most organisms other than Enterobacterales and Acinetobacter have not reached the statistically significant number of 30 isolates. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



INTENSIVE CARE UNIT ANTIBIOGRAM GRAM POSITIVE

Percentage susceptibility gram positive organism 2023

Location	Specimen type	Organism	Number of patients	NITROFURANTOIN	CEFOXITIN	OXACILLIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	MOXIFLOXACIN	CLINDAMYCIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	TETRACYCLINE	Sulfa/Trimeth	TIGECYCLINE	DAPTOMYCIN	LINEZOLID	TEICoplanin	VANCOMYCIN	
SSICU	Blood	Staphylococcus haemolyticus	40	97	8	8	5	8	8		40	21		3	68	53		92	90	95	100	
		Staphylococcus epidermidis	25	96	4	4	0	24	24		52	56		16	64	52		92	96	92	100	
		Staphylococcus hominis	17	87	21	20	13	40	40		80	93		7	67	67		93	93	80	100	
		Staphylococcus aureus	15	100	43	43	7	7	7		86	57		36	86	43		100	100	100	100	
		Enterococcus faecium	12	8			0	0	0				100	0	0				92	75	75	
		Staphylococcus capitis	7	100	86	86	29	100	100		100	100		86	86	100		86	100	100	100	
	Urine	Enterococcus faecium	9	0			0	0	0				100	0	11				78	100	67	
	Pus	Enterococcus faecium	9				0			0		0		100	0			100			100	100
		Staphylococcus aureus	8			38	0			100	100	0			100	63		100			100	100
	Respiratory	Staphylococcus aureus	6	100	67	67	33	0	0		100	100		33	83	33		100	100	100	100	100
	Cerebrospinal fluid	Staphylococcus haemolyticus	2	100	0	0	0	0	0		0	0		0	50	100		100	100	100	100	100
		Staphylococcus hominis	1	100	0	0	0	0	0		0	100		0	0	0		100	100	100	100	100
		Staphylococcus cohnii	1	0	0	0	0	0	0		0	100		0	100	100		100	100	0	100	100

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Except Staphylococcus haemolyticus all other isolates have not reached the statistically significant number of 30 isolates. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern. Staphylococcus species other than S aureus are considered collection contamination/skin flora.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwar Medical Director
--	---	---	---




Candida isolated in ICU

Location	Specimen type	Organism	Number of patients	FLUCONAZOLE	VORICONAZOLE	AMPHO B	CASPOFUNGIN	MICAFUNGIN
SSICU	Blood	Candida albicans	13	62	77	31	100	92
		Candida parapsilosis	8	43	57	57	100	100
		Candida tropicalis	8	29	75	75	100	100
		Candida glabrata	1		0	100	100	100
	Pus	Candida tropicalis	3	100	100	100	100	100
		Candida parapsilosis	1	0	0	100	100	100
	Respiratory	Candida tropicalis	1	100	100	100	100	100
	Urine	Candida tropicalis	6	60	80	80	100	100
		Candida parapsilosis	1	0	0	100	100	100
		Candida dubliniensis	1	0	0	100		
		Candida albicans	5	80	80	40	100	100

Please note: Individual isolates are less than 30 in number, not statistically significant. Antibiogram has been shown with available number of organisms.

Skin flora/Collection contamination in blood culture ICU



Year	No of Contaminants isolated
2022	187
2023	100






Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



SPECIMEN TYPE	OVERALL NO. OF ISOLATES
BLOOD	166
PUS	56
RESPIRATORY	44
URINE	206
CSF	08

MEDICINE

TOP 10 ISOLATES BY SAMPLE



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Dwivedi Medical Director
--	--	---	---



MEDICINE ANTIBIOGRAM GRAM NEGATIVE

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	LEVOFLOXACIN	NORFLOXACIN	CIPROFLOXACIN	OFLOXACIN	NITROFURANTOIN	COTRIAMOXAZOLE	AMOX/CLAV	CEFEPIME	CEF/SUL	CEFUROXIME	CEFTAZIDIME	CEFTRIAAXONE	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMIPENEM	MINOCYCLINE	TIGECYCLINE	FOSFOMYON	COLISTIN	
Medicine	Blood	Escherichia coli	26			4			50	62	23	69	19		19	69	92	89	89	89	89			100	100	
	Urine	Escherichia coli	102		67	11	67	100	49	53	38	71	15	67	24	71	83	71	85	86	86				99	100
		Klebsiella pneumoniae	31	100	0	19	0	0	45	27	23	32	17	0	13	32	39	52	37	33	33	100			61	100
	Respiratory	Klebsiella pneumoniae	8			50			63	63	75	75	50		63	75	75	75	75	75	75		75	75	88	
		Escherichia coli	5			0			60	20	40	20	20		20	20	60	60	60	60	60				100	100
	Pus	Escherichia coli	13			0			25	50	42	58	17		17	58	83	92	67	64	67				100	100
		Acinetobacter baumannii	5			0			20		0	0				0	0	0	0	0	0					100

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Most organisms other than Enterobacterales have not reached the statistically significant number of 30 isolates. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



MEDICINE ANTIBIOGRAM GRAM POSITIVE

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	Oxacillin	Cefoxitin	Levofloxacin	Ciprofloxacin	Cotrimoxazole	Nitrofurantoin	Penicillin	Gentamicin	Gentamicin High Level	Erythromycin	Clindamycin	Tetracycline	Tigecycline	Linezolid	Daptomycin	Teicoplanin	Vancomycin	
Medicine	Blood	Staphylococcus haemolyticus	17	6	0	0	0	40		0	33		13	47	80		87	94	93	93	
		Staphylococcus epidermidis	13	23	23	23	23	69		8	69		23	46	69		100	100	92	100	
		Staphylococcus hominis	13	25	27	33	33	42		8	75		0	58	67		92	83	83	92	
		Staphylococcus aureus	6	33	33	17	17	67		0	50		33	83	83		100	100	100	83	
	Urine	Enterococcus faecium	13			0	0		8	0		100	0		0		92		83	77	
		Enterococcus faecalis	10			10	10		90	100		100	0		0		100	90	100	100	
		Staphylococcus aureus	5	60	60	0	0	100	100	20	100		40	80	100		100	80	100	80	
	Pus	Staphylococcus aureus	10	30	30	10	20	40		0	100		20	80	100		100	100	90	100	100
	Fluid	Staphylococcus haemolyticus	5	0	0	20	20	60		0	40		0	40	60		80	100	100	100	100

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Please note: Individual isolates are less than 30 in number, not statistically significant. Antibiogram has been shown with available number of organisms.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--




CANDIDA ISOLATED IN MEDICINE

Location	Specimen type	Organism	Number of patients	FLUCONAZOLE	VORICONAZOLE	CASPOFUNGIN	MICAFUNGIN	AMPHO B
Medicine	Blood	Candida parapsilosis	9	67	78	100	100	89
		Candida albicans	2	100	100	50	50	100
		Candida tropicalis	1	0	0	100	100	0
	Urine	Candida tropicalis	4	100	100	100	100	100
		Candida albicans	2	100	100	100	100	0
	Respiratory	Candida albicans	3	100	100	100	100	33
		Candida tropicalis	1	0	100	100	100	100
	Fluid	Candida albicans	3	100	100	100	100	67
	Wound	Candida parapsilosis	1	0	100	100	100	100

Candida

Please note: Individual isolates are less than 30 in number, Not statistically significant. Antibiogram has been shown with available number of organisms.

Skin flora/Collection contamination in blood culture








Year	No of Contaminants isolated
2022	46
2023	51

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



**SURGERY :
TOP 10 ISOLATES BY SAMPLE**

SPECIMEN TYPE	OVERALL NO. OF ISOLATES
BLOOD	26
WOUND	297
TISSUE	65
URINE	103
FLUID	21



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	--	--



SURGERY ANTIBIOGRAM GRAM NEGATIVE

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	CIPROFLOXACIN	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	FOSFOMYCIN	COTRIMOXAZOLE	NITROFURANTOIN	AMPICILLIN	AMOX/CLAV	CEFEPIME	CEFUROXIME	CEFTAZIDIME	CEFTRIAZONE	CEF/SUL	PIPTAZ	AMIKACIN	GENTAMYCIN	MEROPENEM	IMPENEM	ERTAPENEM	MINOCYCLINE	TIGECYCLINE	COLISTIN		
Surgery	Blood	Escherichia coli	7	14					86			57	14	0		0	0	86	100	71	100	100				100		
	Wound	Escherichia coli	69	4						42			49	29	12		17	0	64	88	67	84	84	81				100
		Klebsiella pneumoniae	38	40	0					61			51	53	38	0	46	0	53	66	66	63	61	65	100	58		100
		Pseudomonas aeruginosa	26	65	69									77		69		0	65	81	81	73	77					100
		Acinetobacter baumannii	10	0						44				0			0	0	0	0	0	0	0					100
	Tissue	Escherichia coli	15	0						33			47	20	0		0	0	60	73	80	80	80	73				100
		Klebsiella pneumoniae	8	25						63			50	50	25		38	0	50	50	50	50	50	50		63		100
		Acinetobacter baumannii	5	20						40				20			0	0	20	20	20	20	20					100
	Urine	Escherichia coli	51	6		50	50	100	53	50	0	43	32	9	25	24	0	62	84	73	85	85	84					100
		Klebsiella pneumoniae	26	23		100	100	77	46	50		27	21	17	100	27	0	31	46	42	42	38	46					100
	Fluid	Escherichia coli	5	0					60			80	20	0		0	0	80	100	40	80	80	80					100
	Other	Escherichia coli	5	0					20			40	20	20		20	0	60	80	40	60	60	60					100

265

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Most organisms other than Enterobacteriales have not reached the statistically significant number of 30 isolates. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



SURGERY ANTIBIOGRAM GRAM POSITIVE

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	Antibiotics												Reserved / Restricted Drugs				
				COTRIAMOXAZOLE	NITROFURANTOIN	CEFOXITIN	OXACILLIN	PENICILLIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	CLINDAMYCIN	CIPROFLOXACIN	LEVOFLOXACIN	TIGECYCLINE	TETRACYCLINE	DAPTOMYCIN	LINEZOLID	TECOPLANIN	VANCOMYCIN
Surg	Wound	Staphylococcus aureus	50	66	100	34	34	8	78		40	84	10	10	100	96	100	100	100	100
		Staphylococcus epidermidis	14	93	100	0	0	0	93		29	71	36	36	100	71	100	100	100	100
		Enterococcus faecalis	7		100			71		100	0		43	43	100	29	57	100	100	100
		Staphylococcus haemolyticus	6	33	100	0	0	0	33		0	17	0	0	100	100	83	83	83	83
		Enterococcus faecium	5		0			0		100	0		0	0	100	0		100	75	75
	Tissue	Staphylococcus aureus	10	80	100	50	50	10	90		30	90	10	20	100	90	90	100	100	90

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Candida isolates

Candida isolated in Surgery

Location	Specimen type	Organism	Number of patients	Antifungals				
				FLUCONAZOLE	VORICONAZOLE	AMPHO B	CASPOFUNGIN	MICAFUNGIN
Surg	Blood	Candida albicans	2	50	100	0	100	100
		Candida guilliermondii	1	0	0	100	100	100
	Wound	Candida albicans	2	100	100	50	100	100
	Urine	Candida tropicalis	2	100	100	100	100	100
	Fluid	Candida parapsilosis	1	0	100	100	100	100
		Candida tropicalis	1	100	100	100	100	100





Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



**NEONATAL INTENSIVE CARE UNIT:
TOP 10 ISOLATES SAMPLEWISE**

SPECIMEN TYPE	OVERALL NO. ISOLATES
BLOOD	68
PUS	19
URINE	20
ENDOTRACHEAL ASPIRATE	4



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



NICU ANTIBIOGRAM GRAM NEGATIVE

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	COTRIAMOXA ZOLE	NITROFURANTOIN	OFLOXACIN	NORFLOXACIN	LEVOFLOXACIN	CIPROFLOXACIN	FOSFOMYCIN	AMOX/CLAV	CEFEPIME	CEFUROXIME	CEF/SUL	CEFTRIAXONE	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMIPENEM	MINOCYCLINE	COLISTIN
NICU	Blood	Klebsiella pneumoniae	11	100					80	100	82	80	70	0	80	80	100	100	80	90	90		100
		Enterobacter cloacae	6	83					67	100	33	67	33	0	50	67	100	67	100	100	100		100
		Sphingomonas paucimobilis	4	100				100	67			100		0	100	100	33	67	100		100	100	
	Urine	Escherichia coli	10	90	100	67	67		30	100	60	57	14	0	30	90	90	80	100	100	100		100
		Klebsiella pneumoniae	4	50					0	100	25	50	0	0	25	75	75	75	75	75	75		100

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibioqram has been shown with available number of organisms.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



NICU ANTIBIOGRAM GRAM POSITIVE

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	OXACILLIN	CEFOXITIN	COTRIAMOXAZOLE	NITROFURANTOIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	ERYTHROMYCIN	CLINDAMYCIN	GENTAMICIN	TETRACYCLINE	TIGECYCLINE	DAPTOMYCIN	LINEZOLID	TEICOP LANIN	VANCOMYCIN
NICU	Blood	Staphylococcus epidermidis	9	11	11	22	100	0	11	11	11	22	22	89		100	100	100	100
		Staphylococcus haemolyticus	6	0	0	50	100	0	0	0	0	33	0	100		100	100	100	83
		Staphylococcus hominis	4	25	25	50	100	0	50	50	0	25	100	75		100	100	100	100
	Wound	Staphylococcus aureus	10	20	20	90	100	10	10	10	30	100	90	100	100	100	100	100	100

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Candida

Candida isolated in NICU

Location	Specimen type	Organism	Number of patients	Fluconazole	Voriconazole	Ampho B	Caspofungin	Micafungin
NICU	Blood	Candida albicans	2	100	50	0	100	100
		Candida parapsilosis	1	0			100	100
		Candida tropicalis	1	100	100	100	100	100
	Cerebrospinal fluid	Candida albicans	2	50	100	50	100	100
	Fluid	Candida parapsilosis	1	100	100	100	100	100



Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	---	--



**OBSTETRICS AND GYNAECOLOGY:
TOP 10 ISOLATES SAMPLEWISE**

SPECIMEN TYPE	OVERALL NO OF ISOLATES
WOUND	20
URINE	28



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	--	--



OBSTETRICS AND GYNECOLOGY ANTIBIOGRAM GRAM NEGATIVE

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	COTRIMOXAZOLE	NITROFURANTOIN	CIPROFLOXACIN	NORFLOXACIN	OFOXACIN	LEVOFLOXACIN	FOSFOMYCIN	AMOX/CLAV	CEFUROXIME	CEF/SUL	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMPENEM	TIGECYCLINE	COLISTIN		
Obgyn	Blood	Escherichia coli	2	100		0				100	100	0	0		0	0	100	100	100	100	100	100	100	100		
	Urine	Escherichia coli	22	59	100	27	56	44		96	73	8	0	56	18	46	86	91	86	92	91	92			100	
		Klebsiella pneumoniae	3	100	0	100	100	100		100	67	100	0	100	67	100	100	100	100	100	100	100			100	
		Pseudomonas aeruginosa	1			0				0				0	0		0	0	0	0	0				0	
	Wound	Escherichia coli	8	50		0					25	0	0		0	13	63	75	63	88	88	88			100	
		Klebsiella pneumoniae	1	100		0					100	0	0		0	0	100	100	0	100	100	100	100			100
		Achromobacter xylosoxidans	1	100		0				0			0	100		0	100	0	0	100		100				
		Citrobacter koseri	1	100	100						100	0	0		100	100	100	100	100	100	100	100			100	
	Other	Escherichia coli	4	25		0					75	0	0		0	0	100	75	75	100	100	100			100	
		Klebsiella pneumoniae	2	50		50						50	50	0		50	50	50	100	50	100	100	50		100	

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Please note: Other than E. coli individual isolates are less than 30 in number. Not statistically significant numbers.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



OBSTETRICS AND GYNECOLOGY ANTIBIOGRAM GRAM POSITIVE

No significant number of Gram Positive isolates in 2023

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	NITROFURANTOIN	COTRIAMOXAZOLE	OXACILLIN	CEFOXITIN	LEVOFLOXACIN	CIPROFLOXACIN	RIFAXIMIN	PENICILLIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	CLINDAMYCIN	TETRACYCLINE	TEICoplanin	VANCOMYCIN	DAPTOMYCIN	LINEZOLID	TIGECYCLINE	
Obgyn	Blood	Staphylococcus hominis	1	100	0	100	100	0	0	100	100	100		0	0	100	100	100	100	100		
		Streptococcus agalactiae	1	100				100			100						0		100	100		
		Enterococcus faecium	1	0				0	0		0		100		0	0	0	0	0	0	100	
		Staphylococcus, coagulase negative	1	100	100	100	100	100	100	100	100	0	100		100	100	100	100	100	100	100	
	Urine	Streptococcus agalactiae	1	100				100			100						0		100	100		
		Enterococcus faecium	1	0				0	0		0		100		0	0	0	100	100	100		
	Other	Staphylococcus haemolyticus	2	100	50	50	50	50	0	100	50	100			50	100	100	100	100	100	100	100
		Enterococcus faecium	1	0				0	0		0		100		0	0	0	0	0	100	100	
		Enterococcus faecalis	2	100				50	50		100		100		0	0	0	100	100	0	50	100
		Staphylococcus aureus	1	100	100	100	100	100	100	100	100	0	100		0	100	100	100	100	100	100	100

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic





Please note: Individual isolate numbers are in single digits. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



**ORTHOPAEDICS :
TOP 10 ISOLATES SAMPLEWISE**

SPECIMEN TYPE	OVERALL NO OF ISOLATES
WOUND	77
TISSUE	46
URINE	42
BLOOD	22



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	--	--



ORTHO PAEDICS ANTIBIOGRAM GRAM NEGATIVE

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	Antibiotics												Other Antibiotics					
				COTRIAMOXAZOLE	CIPROFLOXACIN	LEVOFLOXACIN	FOSFOMYCIN	AMOX/CLAV	CEFUROXIME	CEF/SUL	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMPENEM	TIGECYCLINE	COLISTIN
ortho	Urine	Escherichia coli	23	22	4		96	13	0	0		4	14	38	48	44	39	39	46		100
		Klebsiella pneumoniae	11	46	27		82	36	27	0		18	36	46	46	55	46	46	46		100
	Wound	Escherichia coli	13	54	8		100	39	8	0		15	39	54	100	92	77	77	77		100
		Pseudomonas aeruginosa	8		75	75				0	63		63	63	88	75	75		88		100
		Acinetobacter baumannii	7	43	0					0		0	0	0	0	0	0		0		100
		Klebsiella pneumoniae	7	57	14		71	29	14			14	14	14	57	57	43	43	43	57	100
	Tissue	Escherichia coli	11	27	0		100	27	0	0		0	0	36	100	91	55	55	55		100
		Pseudomonas aeruginosa	8		75	63				0	75		75	63	88	88	63		75		100

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Please note: Individual isolates are less than 30 in number. Not statistically significant. AntibioGram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



ORTHOPAEDICS ANTIBIOGRAM GRAM POSITIVE

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	Antibiotics										Reserved / Restricted Drugs						
				COTRIAMOXAZOLE	NITROFURANTOIN	OXACILLIN	CEFOXITIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	CLINDAMYCIN	TETRACYCLINE	DAPTOMYCIN	LINEZOLID	TIGECYCLINE	TEICoplanin	VANCOMYCIN	
ortho	Wound	Staphylococcus aureus	23	74	100	39	39	9	13	13	78	48	96	96	96	100	100	100	100	100
	Tissue	Staphylococcus aureus	12	58	100	50	50	17	17	25	75	42	100	92	100	100	100	100	100	100
	Other	Staphylococcus aureus	5	60	100	60	60	0	20	20	40	60	100	100	100	100	100	100	100	100

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic





Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



PEDIATRICS& PICU : TOP 10 ISOLATES SAMPLEWISE

SPECIMEN TYPE	OVERALL NO OF ISOLATES
BLOOD	36
WOUND	31
URINE	30



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	--	--



Paediatrics Antibiogram

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	COTRIMOXAZOLE	CIPROFLOXACIN	LEVOFLOXACIN	FOSFOMYCIN	AMOX/CLAV	AZTREONAM	CEFEPIME	CEF/SUL	CEFTRIAZONE	CEFUROXIME	PIPTAZ	AMIKACIN	GENTAMICIN	IMIPENEM	MEROPENEM	ERTAPENEM	MINOCYCLINE	COLISTIN
Paed	Blood	Salmonella Typhi	5	100	0	0		100		100		100		100			100	100	100		
	Wound	Escherichia coli	6	0	0	0		50	0	33	17	25	25	17	100	80	50	60	50	100	100
		Klebsiella pneumoniae	2	50	0			50		50	100	0	0	100	100	100	50	100	100		100
	Urine	Escherichia coli	20	37	5		100	20		47	47	32	16	47	90	63	74	68	68		100
		Klebsiella pneumoniae	4	25	0		50	0		0	0	0	0	0	25	0	0	25	25		100
	Respiratory	Pseudomonas aeruginosa	2		100	100					100	100			100	0	0	0			100

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--




Please note: Individual isolates are less than 30 in number. Antibiogram has been shown with available number of organisms.

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	CEFOXITIN	OXACILLIN	COTRIAMOXAZOLE	NITROFURANTOIN	CIPROFLOXACIN	LEVOFLOXACIN	PENICILLIN	GENTAMICIN	ERYTHROMYCIN	CLINDAMYCIN	TETRACYCLINE	TIGECYCLINE	LINEZOLID	DAPTOMYCIN	TEICoplanin	VANCOMYCIN
Paed	Blood	Staphylococcus aureus	2	0	0	0	100	0	0	0	100	0	100	50		100	100	100	100
		Staphylococcus hominis	3	33	33	100	100	100	100	33	67	33	67	100		67	67	67	67
		Staphylococcus epidermidis	5	0	0	80	100	60	60	0	60	20	40	100		100	100	60	100
	Wound	Staphylococcus aureus	14	50	50	50	100	0	0	0	93	43	86	100	100	100	93	100	93

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Skin flora/Collection contamination in blood culture – Paediatrics & PICU



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--






ONCOLOGY

Please note: Individual isolates are less than 30 in number. Antibigram has been shown with available number of organisms.

Please prescribe empiric therapy as per policy guidelines outlined in the section on therapy of common conditions and Febrile neutropenia

SPECIMEN TYPE	NO OF ISOLATES
BLOOD	42
PUS/TISSUE	19
URINE	10



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



ONCOLOGY

Percentage Susceptibility Gram Negative Organism 2023

Location	Specimen type	Organism	Number of patients	COTRIAMOXAZOLE	CIPROFLOXACIN	LEVOFLOXACIN	AMOX/CLAV	CEFEPIME	CEF/SUL	CEFTRIAZONE	CEFUROXIME	CEFTAZIDIME	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMIPENEM	TIGECYCLINE	MINOCYCLINE	FOSFOMYCIN	COLISTIN		
ONCO	Blood	Enterobacter cloacae	5	40	40		0	40	0	20	20		40	40	40	40	40	40			20	40		
		Pseudomonas aeruginosa	4		25	25		50	0				50	50	100	100	50		50				25	
		Pseudomonas stutzeri	3	0	0	0		100	0				100	100	100	100	100	100						
		Acinetobacter baumannii	2	50	50			50	0	50				50	50	50	50	50					100	
		Acinetobacter junii	2	50	100	100		100	0	0			0	0	0	100	0	0	0	100			0	
		Klebsiella pneumoniae	2	0	0		0	0	0	0	0	0		0	50	50	50	50	50			100	0	
		Escherichia coli	1	100	0		100	0	0	0	0	0		100	100	100	100	100	100			100	100	
		Pseudomonas alcaligenes	1	0	0	0		100	0				100		100	0	100	100						
		Proteus mirabilis	1	0	0		0	0	0	0	0	0		100	0	0	0	0	0				100	
		Aeromonas hydrophila	1	100	100	100		100	0				100	0	100	100	0	0	0					
	Salmonella enterica	1	100	100		100	100	0	100	100			100	100	100	100	100	100			100	100		
	Tissue	Klebsiella pneumoniae	1	0	0		0	0	0	0	0		0	100	100	0	0	0	100			100	100	
		Citrobacter freundii	1	0	0		100	0	0	0	0		0	100	0	0	0	100	0			100	0	
		Escherichia coli	1	0	0		0	0	0	0	0		0	0	0	0	0	0	0			100	100	
	Wound	Klebsiella pneumoniae	2	50	0		0	50	0	0	0		50	50	50	50	50	50	0	50			100	100
		Pseudomonas aeruginosa	2		100	50		0	0				0	0	100	100	100	100					0	
		Escherichia coli	1	0	0		0	0	0	0	0		0	100	100	100	100	100					100	100
		Proteus mirabilis	1	100	0		100	100	0	100	100		100	100	100	100	100	100	0				100	
		Pseudomonas stutzeri	1	0	0	100		100	0				100	100	100	0	0	0	0					
	Urine	Escherichia coli	4	0	0		0	25	0	0	0		50	100	75	75	100	100	100			100	100	
Klebsiella pneumoniae		2	0	0		50	50	0	0	0		50	100	100	100	100	100	100			50	100		

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibigram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	---	--



ONCOLOGY GRAM NEGATIVE

Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	COTRIAMOXAZOLE	NITROFURANTOIN	OXACILLIN	CEFOXITIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	CLINDAMYCIN	TETRACYCLINE	DAPTOMYCIN	LINEZOLID	TIGECYCLINE	VANCOMYCIN	TEICOPLANIN	
onco	Blood	Staphylococcus aureus	3	100		33	33	0	0	0	100		33	100	100	100	100		100	100	
		Staphylococcus epidermidis	1	0		0	0	0	0	0	0			0	0	100				100	100
		Staphylococcus hominis	3	50		33	0	0	0	0	0	100		0	50	0	67	50		100	100
	Wound	Staphylococcus aureus	5	100		20	20	0	20	20	100		40	100	100	100	100	100	100	100	100
		Staphylococcus haemolyticus	2	50		0	0	0	0	0	0			0	50	100	100	50	100	100	100
		Staphylococcus hominis	1			100											100		100		
	Urine	Enterococcus faecium	2		0			0	0	0		100	0		0		100		50	50	

Reserved / Restricted Drugs :Not to be used empirically unless justified
 Will be useful clinically
 Will be useful clinically in about 2/3 cases
 Will not be useful clinically
 Not tested/ not appropriate antibiotic

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



Skin flora/Collection contamination in blood culture - Oncology

ONCOLOGY GRAM POSITIVE

Please note: Individual isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Percentage Susceptibility Gram Positive Organism 2023

Location	Specimen type	Organism	Number of patients	Antibiotics												Reserved / Restricted Drugs																								
				COTRIAMOXAZOLE	NITROFURANTOIN	OXACILLIN	CEFOXITIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	CLINDAMYCIN	TETRACYCLINE	DAPTOMYCIN	LINEZOLID	TIGECYCLINE	VANCOMYCIN	TEICoplanin																				
ONCO	Blood	Staphylococcus aureus	3	100		33	33	0	0	0	100		33	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100				
		Staphylococcus epidermidis	1	0		0	0	0	0	0	0	0	0	0	0	100		0	0	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
		Staphylococcus hominis	3	50		33	0	0	0	0	0	100		0	50	0	67	50		100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
	Wound	Staphylococcus aureus	5	100		20	20	0	20	20	100		40	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
		Staphylococcus haemolyticus	2	50		0	0	0	0	0	0	0	0	0	50	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
		Staphylococcus hominis	1			100																																		
	Urine	Enterococcus faecium	2		0				0	0	0		100	0																										

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--






ALL OPD

SPECIMEN TYPE	NUMBER OF ISOLATES
BLOOD	59
WOUND	362
SPUTUM	12
URINE	515

Blood isolates are from cultures taken in the Emergency Department

Skin flora/Collection contamination in EMD	
Staphylococcus epidermidis	7
Staphylococcus haemolyticus	4
Staphylococcus hominis ss. hominis	3
Kocuria kristinae	1
Kocuria rosea	1
Micrococcus luteus	1
Staphylococcus arlettae	1
Staphylococcus saprophyticus ss. saprophyticus	1
Staphylococcus caprae	1
Streptococcus intermedius	1
Staphylococcus xylosus	1
Total	22

OPD : TOP 10 ISOLATES SAMPLEWISE



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



ALL OPD GRAM NEGATIVE

Percentage susceptibility gram negative organism 2023

Location	Specimen type	Organism	Number of patients	COTRIMOXAZOLE	NITROFRANTOIN	OFLOXACIN	LEVOFLOXACIN	CIPROFLOXACIN	NORFLOXACIN	FOSFOMYCIN	AMOX/CLAV	CEFEPIME	Cefuroxime oral	CEFUROXIME	CEFTAZIDIME	CEFTRIAXONE	CEF/SUL	PIPTAZ	AMIKACIN	GENTAMICIN	MEROPENEM	ERTAPENEM	IMPENEM	TIGECYCLINE	FOSFOMYCIN	COLISTIN	
OPD	Urine	Escherichia coli	286	51	86	36		13	45	99	55	37	15	27	64	28	78	78	81	73	92	92	92		99	100	
		Klebsiella pneumoniae	87	48	33	74	0	29	74	76	38	30	18	41	63	39	45	45	65	64	55	60	56		76	100	
		Pseudomonas aeruginosa	28				31	32					52			48		46	46	52	60	46		48			100
		Enterobacter cloacae	14	79	33	83		57	100	50	7	88	13	50	67	71	71	71	86	93	88	93	88		50	100	
		Proteus mirabilis	11	46		0		27	33	100	82	88	88	100	100	91	100	100	100	82	64	100	100	38		100	
	Wound	Pseudomonas aeruginosa	50				66	69					88			86		79	79	82	71	74		82			100
		Escherichia coli	45	71				4			44	49	13			29	78	78	91	76	93	91	96		100	100	
		Klebsiella pneumoniae	24	58	0	100		42	100		61	55	41	100	100	54	65	65	78	83	86	71	77	65	87	100	
		Enterobacter cloacae	10	90				80			10	90	10			60	90	90	90	100	90	90	90	90	80	100	
		Proteus mirabilis	5	20			0	0			0	40	50			0	50	80	80	40	40	80	75	20		100	
	Other	Escherichia coli	10	40					30			70	40	30			30	80	80	100	90	100	90	100	100	100	

- Reserved / Restricted Drugs :Not to be used empirically unless justified
- Will be useful clinically
- Will be useful clinically in about 2/3 cases
- Will not be useful clinically
- Not tested/ not appropriate antibiotic

Please note: Some isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--



ALL OPD GRAM POSITIVE

Percentage susceptibility gram positive organism 2023

Location	Specimen type	Organism	Number of patients	CEFOXITIN	OXACILLIN	COTRIAMOXAZOLE	NITROFURANTOIN	RIFAMPICIN	PENICILLIN	CIPROFLOXACIN	LEVOFLOXACIN	GENTAMICIN	GENTAMICIN HIGH LEVEL	ERYTHROMYCIN	CLINDAMYCIN	TETRACYCLINE	VANCOMYCIN	TEICoplanin	DAPTOMYCIN	LINEZOLID	TIGECYCLINE	
	Wound	Staphylococcus aureus	96	35	35	61		97	5	11	12	78		48	87	93	100	100	95	99	100	
		Staphylococcus epidermidis	14	29	29	71		79	0	36	43	86		14	57	93	100	100	93	93	100	
		Enterococcus faecalis	10						100	44	44			100	11		0	100	100	67	100	100
	Urine	Enterococcus faecalis	16				94		81	6	6			100	19		0	94	94	69	100	
		Staphylococcus aureus	10	50	50	60	100	100	10	0	0	80		40	100	100		100	100	100	100	
		Enterococcus faecium	8				0		0	0	0			100	14		29	57	57		71	
		Staphylococcus saprophyticus	5	60	60	80	100	100	0	100	100	100			0	20	100	100	100	100	100	

■ Reserved / Restricted Drugs :Not to be used empirically unless justified
■ Will be useful clinically
■ Will be useful clinically in about 2/3 cases
■ Will not be useful clinically
■ Not tested/ not appropriate antibiotic

Please note: Some isolates are less than 30 in number. Not statistically significant. Antibiogram has been shown with available number of organisms. Hence these isolates may not be truly representative of the antimicrobial susceptibility pattern.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	--	--




Index of multidrug resistance

Shows the trend of multidrug resistance of common pathogens for the years 2019-2023

Multi drug resistance index is calculated by formula $\frac{\text{Number of drugs resistant}}{\text{Number of drugs tested}}$

Value closer to 0.2 is considered the best



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Gulwani Medical Director
--	---	---	--



6. Antimicrobial therapy in hospitalized patients

Antimicrobial therapy is used in hospitalized patients in three situations

1. **Empiric therapy** before the causative organism has been identified
2. **Definitive therapy** once the causative organism is identified
3. **Prophylactic therapy** to prevent infection, eg. surgical prophylaxis

Empiric therapy

Patient requiring empirical antimicrobial therapy should be classified into three types (Table 1) depending on the past history, prior exposure to health care, previous antimicrobials and associated co-morbidities. Antimicrobial should then be chosen according to the site of infection and suspected micro-organism based on local hospital microbiologic data (antibiogram). Appropriate cultures must be sent prior to antimicrobial therapy. Identification of the micro-organism will then dictate definitive therapy and also contribute to the hospital antibiogram for choosing empiric therapy.

Please note: Empiric therapy will be started only after appropriate samples for culture have been taken

TABLE 1- Patient Types for selecting empiric antimicrobial therapy

Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
No contact with health care system	Contact with health care system (e.g. recent hospital admission, nursing home, dialysis) without invasive procedure within last 90 days. Current hospitalization less than 7 days	Current hospitalization > 7 days. Invasive procedures within last 90 days
No prior antimicrobial treatment	Recent antimicrobial therapy (within last 90 days)	Recent & multiple antimicrobial therapies within last 90 days Major invasive procedures done
No procedures done	Minimum procedures done	
Patient young with only a few co-morbid conditions.	Patient old with Multiple co-morbidities.	Cystic fibrosis, structural lung disease, advanced AIDS, neutropenia other Severe immunodeficiency

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department by Med Dir	Approved by Dr SK Gulwani Medical Director
--	---	---	--



7. Empiric Antimicrobial choice

Recommended antimicrobials for common conditions are listed below. This guide is a broad outline; not all-inclusive and; not meant to replace treating physician’s judgment.

Table 1: Acute gastroenteritis

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
Acute gastroenteritis	<p>Most cases are self-limited and require only supportive treatment and hydration or being viral in nature. Selected sick patients can be treated as per following guidelines. Tab Cefixime 200mg BD</p> <p>If stool examination shows invasive diarrhoea (> 5 leucocytes /HPF or blood in the stool).</p> <p>Then consider stool culture followed by therapy as per AST</p>		

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Table 2: Pneumonia

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
Pneumonia	<p>1] For non-ICU patients with community acquired pneumonia (CAP)Ceftriaxone (2g IV q24h X 5-7 days)/ Amoxicillin/Clavulanic acid (1.2g q8h IV)</p> <p style="text-align: center;">+</p> <p>Macrolide (Azithromycin- 500mg IV/PO once a day), x 5-7 days).</p>	<p>Late Onset HAP/VAP (For more than 48 hours of hospitalization but less than 7 days)</p> <p>If septic shock or multisystem organ failure, Imipenem 0.5-1 gm q6h or Meropenem 1-2 gm q8h</p>	<p>Late onset HAP/VAP suspected MDR Gram negative –</p> <p>Imipenem (0.5-1 gm q6h /Meropenem (1-2 g IV q8h)</p> <hr/> <p>Suspected XDR Gram negative</p> <p>Colistin 4.5 MU/BD</p> <p>Suspected MRSA-</p> <p>Vancomycin (1g IV q12h OR Teicoplanin (400mg IV q12h for 3 doses, then q24h)</p>
	<p>2] ICU patients with CAP</p> <p>Ceftriaxone (2g IV q24hr X 5-7 days)/</p> <p>Amoxicillin/Clavulanic acid (1.2g q8hr IV)</p> <p style="text-align: center;">+</p> <p>Macrolide: Azithromycin- 500mg IV/PO q24h)/ Doxycycline 100mg PO q12h x 5-7 days).</p> <p>If aspiration is suspected clindamycin 600mg q8h</p> <p>Early onset HAP/VAP</p>		<p>For suspected VRE-</p> <p>Linezolid (600mg IV/PO q12hr)**x 7-14 days</p> <p>For suspected Fungal (Filamentous fungi/mould) infections-</p> <p>Consider Antifungals in Immunocompromised host. Add Liposomal Amphotericin B. Substitute Voriconazole, if Aspergillus suspected on radiological evidence or galactomannan positive</p> <hr/> <p>If PCP suspected-</p> <p>add TMP-SMX or Clindamycin</p>

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



	<p>(less than 48 hours admission) Antimicrobial choice as above unless Pseudomonas or Gram negative bacilli are suspected. Then use Cefoperazone-Sulbactam* (1.5g-3gm q6h) or piperacillin-tazobactam (PIP-Taz) 4.5gm q6h</p>		
H1N1 Flu-like illness	<p>Look for typical viral symptoms such as sneezing and running nose. If fever, sore throat, dry cough and viral symptoms present, initiate Oseltamivir 75 mg BD x 5 d without waiting for confirmation by PCR</p>		
<p>Note :</p> <ol style="list-style-type: none"> 1. Fluoroquinolones should not be used for empiric treatment. 2. Fluoroquinolones should not be used routinely for treating Acute exacerbation of COPD 3. In the uncommon scenario of hypersensitivity to β-lactams, respiratory Fluoroquinolones (e.g. levofloxacin 750 mg daily) may be used if tuberculosis is not a diagnostic consideration at admission. Patients should also undergo sputum testing for acid-fast bacilli simultaneously if fluoroquinolones are being used in place of β-lactams. 4. **Patients with suspected MRSA infection, we recommend the use of empiric Vancomycin or Teicoplanin. The use of linezolid in India should be reserved because of its potential use in extensively drug-resistant tuberculosis. 5. Suspected viral pneumonia [influenza] Oseltamivir and/or Zanamavir should be given. 6. In late HAP/VAP with suspected Acinetobacter infection combination of Colistin + carbapenem / sulbactam. 			

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



7. Duration of treatment for community acquired pneumonia should be minimum 5-7 days and patient should be afebrile 48-72 hours prior to stopping treatment.
8. For ESBL / MRSA health care associated pneumonia minimum duration of treatment should be 10-14 days.
9. For proven pseudomonal / Acinetobacter health care associated pneumonia treatment should be for minimum 2 weeks and preferably combination of antimicrobial therapy should be used.
10. Colonization should be suspected if respiratory secretions culture show growth but following features are absent like Fever, leukocytosis, increased bronchorrhea, increasing oxygen requirement, new lung infiltrates.
11. In presence of Fever, leukocytosis, increased bronchorrhea, increasing oxygen requirement but absence of lung infiltrates with positive cultures [MDR GNB / MRSA] to be treated as health care associated tracheobronchitis with appropriate broad spectrum antimicrobials.
12. Aerosolised Tobramycin/ Colistin can be added to IV antimicrobials as an adjunctive therapy for MDR gram negative infection with specialized nebulisers.

Bharati Hospital and Research Centre

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



Table 3: Meningitis

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
Meningitis	<p>1] Age 2yrs-50yrs Vancomycin 1gm q12h + Ceftriaxone 2gm q12h</p> <p>2] Age > 50yrs Above Antimicrobials + Ampicillin 2gm q4h</p>	Vancomycin 1gm q12h + cefepime 2gm q12h /Ceftazidime 2gm q8h	<p>Empirical Therapy Vancomycin 1gm q12h + Colistin 4.5 MU BD+/- Meropenem 2gm q8h. Consider Intrathecal Gentamicin/ Colistin 4.5 MU BD</p> <p>Organism specific A] Suspected MRSA Meningitis – Vancomycin 1gm q12h +/- Rifampicin 600mg q12hor Linezolid 600mg q12h</p> <p>B] ESBL Gram negative/Pseudomonas or Acinetobacter (MDR / XDR) Meropenem 2gm q8h + Colistin 4.5 MU BD .</p>

Note : Intrathecal/ Intraventricular route dosage-

Vancomycin 10-20mgq24h;

Gentamicin 4-8 mgq24h;

Amikacin 30-50mg q24h;

Colistin 5-20mg q24h[1mg = 12,500 units]

IV Dexamethasone should be given in suspected pneumococcal meningitis before antimicrobial therapy and should be continued only if Gram stain / Culture confirms pneumococcal etiology

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Table 4: Urinary tract infection

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
UTI	<p>Asymptomatic bacteriuria No empirical therapy. Send C/S.</p> <p>Non complicated UTI (Cystitis, Urethritis, No evidence of obstructive uropathy) PO TMP SMX 160/800 q12h / PO Nitrofurantoin 100 mg q12h</p> <p>Acute Uncomplicated Pyelonephritis Fluroquinolones Ofloxacin 400 mg q12h OR Gentamicin 3 – 5 mg/kg q24h 5-7 days If hospitalized Ceftriaxone 1gm q12h</p>	<p>Complicated UTI (Obstruction, reflux, azotemia, CAUTI) IV Meropenem 1gm q8h/ IV Imipenem-cilastatin 0.5 gm q6h</p> <p>Complicated Pyelonephritis (Obstruction, reflux, azotemia, CAUTI, Shock, perinephric abscess) Meropenem 1gm q8h/Imipenem-cilastatin 0.5 gm q6h in presence of obstruction</p>	<p>Complicated UTI and Pyelonephritis (Suspected MDRO's/ Post renal transplant/ Recurrent UTI's) IV Meropenem 1gm q8h/IV Imipenem-cilastatin 500mg q6h +/- IV Colistin 4.5 MUBD</p> <p>If MRSA or enterococcus, Consider Vancomycin 1 gm q12h/ Teicoplanin 400 mg q24h</p> <p>look for obstruction. Surgical management is mandatory to relieve obstruction</p>

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Lower Urinary tract infection(UTI) in antenatal patients up to 20 weeks gestation	OPD- Cap.Amoxyicillin500 mg q8h PO In-patient IV Ceftriaxone 1gm q12h		Meropenem 1gm q8h Or Colistin 4.5MUBD
Lower Urinary infection(UTI) in antenatal patients after 20 weeks gestation	OPD Tab. Nitrofurantoin SR100 mg BD oral/ Oral Fosfomycin 1g sachet single dose/Oral Cefixime 200mg BD x 3-5 days Inpatient: Ceftriaxone 1gm q12h	IV PIP-TZ 4.5 gm q6h	Meropenem 1gm q8h

Table 4 a : Pelvic Inflammatory Disease

Pelvic inflammatory disease: Mild to moderate	Tab Cefixime Plus Tab Metronidazole Plus Cap Doxycycline	Levofloxacin Plus Metronidazole or Ceftriaxone	
Pelvic inflammatory disease: Severe	Clindamycin Plus Gentamicin	IV PIP-TZ 4.5 gm q6h or Imipenem	

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



Table 5- Skin & soft tissue infections

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
Erysipelas / uncomplicated cellulitis	IV Ceftriaxone 2 gm q24h If beta lactam allergy IV Clindamycin 600 – 900 mg q8h		
Necrotizing infection of skin/fascia and muscle	IV Ceftriaxone 2gm q12h + IV Clindamycin 600-900mg q8h / IV Metronidazole 500mg q6h If Suspected MRSA IV Vancomycin 1 gm q12h/ IV Teicoplanin 400 mg q24h		
Fournier gangrene	Mixed aerobic and anaerobic cover including S.aureus MRSA cover IV Vancomycin 1gm q12h If pseudomonas suspected IV PIP-TZ 4.5gm q6h		
Diabetic foot	IV Co-amoxiclav 1.2 gm q8h if beta lactam allergy- IV Clindamycin 600 q8h	IV PIP-TZ 4.5 gm q6h If Suspected MRSA infection IV Vancomycin 1 gm q12h	IV Meropenem 1gm q8h or IV Imipenem-Cilastatin 1gm q6h. If MRSA suspected IV Vancomycin 1gm q12h

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
---	--	--	---



Table 6- Bone and joint infections

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
Acute Osteomyelitis / Septic Arthritis	Ceftriaxone IV q12h OR Co-amoxiclav 1.2 gm q8h with/without Gentamicin 3 –5 mg/kg q24h If MRSA suspected- Vancomycin 1gm IV q12h	-	-
Early implant associated infection (< 3 months)	-	Usual Suspected organism- Staph aureus/ MRSA IV Vancomycin 1 gm q12h/ Teicoplanin(400mg IV q12h for 3 doses, then q24h) + If Suspected MDR Gram negative organism IV Meropenem 1gm q8h IV Imipenem + Cilastatin 1gm q6h IV/IV Colistin	-
Late implant associated infection (after 3 months)	-	-	Usually low grade infection If Coagulase negative staphylococcus suspected - IV Vancomycin 1 gm q12h / Teicoplanin (400mg IV q12h for 3 doses, then q24h) If Anaerobe (Propionibacterium acne) suspected IV Clindamycin 600-900 mg q8h.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



Table 7 Intra-abdominal infections -

Name of condition	Patient Type 1(Community acquired)	Patient Type 2 (Healthcare associated)	Patient Type 3 (Nosocomial Infections)
A) Extra – biliary	IV Ceftriaxone1-2 gm q12h+IV Metronidazole500mg q8h or IV PIP-TZ 4.5gm q6h	IV Meropenem 1gm q8h/ IV Imipenem-cilastatin 500mg q6h	IV Meropenem 1gm q8h IV Imipenem - cilastatin500mg q6h In case of suspected Acinetobacter or XDR Gram negative organisms Colistin 4.5 MU BD If MRSA or Enterococcus suspected IV Vancomycin 1 gm q12h / Teicoplanin(400mg IV q12h for 3 doses, then q24h) If VRE suspected Linezolid 600 mg IV q12h suspected, Add Fluconazole 400 mg IV q24h If non albicans Candida- IV Caspofungin 70 mg stat and 50 mg q24h or Ampho B
B)Intra Abdominal Biliary	IV Ceftriaxone1-2 gm q12h + IV Metronidazole500mg q8h or IV PIP-TZ 4.5gm q6h	IV Meropenem 1gm q8h / IV Imipenem- cilastatin 500mg q6h	Eg- Acute cholangitis following bilio enteric anastomosis IV Meropenem 1gm q8h/ IV Imipenem - cilastatin500mg q6h .If MRSA or Enterococcus suspected IV Vancomycin 1 gm q12h / Teicoplanin(400mg IV q12h for 3 doses, then q24h). If VRE suspected Linezolid 600 mg IV q12h If Fungal Infection suspected, Add Fluconazole 400 mg IV q24h If non Albicans Candida IV Caspofungin 70 mg stat and 50 mg q24h Or Ampho B

Notes: Metronidazole dosing based on Pharmacokinetic studies is 1.5 GI q24h. 50 mg q24h Or Ampho B

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Tilwahi Medical Director
--	---	--	--



Table 8: Infective Endocarditis

Name of condition		
Native Valve Endocarditis	IV Ceftriaxone	Alternative Penicillin G2-3mu IV q4h or Vancomycin500 mg q12h for 4weeks Ceftriaxone 2 gmq24h for 2 weeks plus Gentamicin 3mg per kg divided into equal doses q8h for 2 weeks
Prosthetic valve endocarditis	Cloxacillin 2gm IV q4h for 4-6 weeks or IV Vancomycin500 mg q12h for 4-6 weeks	IV Cefazolin 2g q8h
<p>Note:- If Penicillin resistant Streptococci - Ceftriaxone 2 gram per day IV q24h for 6 weeks plus Gentamicin 3mg per kg divided into equal doses q8h for 6 weeks Enterococci – Ampicillin 2gm IV q4h + Gentamicin3mg per kg divided into equal doses q8hboth 4-6 weeks or Vancomycin 500 mg q12h + Gentamicin for 4weeks. Staphylococci –Nafcellin or Oxacillin 2gm IV 4 hourly for 4-6 weeks or Vancomycin 15 mg /kg IV 12 hourly for 4-6 weeks</p>		

Bharati Hospital and Research Centre

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	---	--	--



**Table 9: Malaria, Leptospirosis, Scrub Typhus, Enteric fever
(IN LABORATORY CONFIRMED CASES)**

Plasmodium Vivax Malaria	Chloroquine Sensitive	Chloroquine resistant –
	Chloroquine (10mg base/kg stat followed by 5 mg/kg at 12,24,36 hours) plus Primaquine (7.5 mg (base) q12h PO x14days) (Primaquine should not be given in severe G6PD deficiency)	any of the ACT therapy excluding SP 1. Artesunate +Amodiaquine 2. Artesunate +Mefloquine 3. Dihydroartemisin plus piperazine
Plasmodium Falciparum Malaria	OPD Artesunate(2.4 mg/kg at 12 & 24 hours) plus Sulfadoxine (25 mg/ kg) &Pyrimethamine (1.25 mg/kg) as a single dose or Artesunate(same dose as above) plus Amodiaquine(10mg) base per kg OD for 3 days (Fixed dose combinations are available) or Artemether plus Lumefantrine(1.5/9mg/kg BD for 3 days) Drug combination of A+L(mg)available 40+240:60+360:80+480 or Artesunate +Mefloquine (25mg base/kg –total) (8mg/kg once a day for 3 days)	Drug resistant Falciparum Malaria Artesunate 2.4 mg/kg for 7 days or Quinine (10mg/kg TDS for 7 days plus one of the following three 1. Tetracycline 4mg/kg Odx7 days 2. Doxycycline 3mg/kg OD x 7days 3. Clindamycin 10mg/kg BD x 7days
	Hospitalized patient Artesunate IV 2.4 mg/kg at 12 & 24 hours and 2.4 mg/kg q24h X 5 days + Doxycycline 100mg q12h x 7 days	
Leptospirosis (Mild)	Doxycycline 100mg q12h x 7 days	Alternative: Amoxicillin (500 mg)PO TDS x7 days Ampicillin (500mg)PO TDS x 7 days
Leptospirosis (Moderate or Severe)	Ceftriaxone (1gm 12 hourly x7 days or Cefotaxime (1gm 6 hourly IV x 7 days	Alternative Penicillin (1.5 million units /IV /IM 6 hourly x7 days
Scrub Typhus	Doxycycline (100mg) BD x 7 to 15 days or Azithromycin (500mg) OD x 3days	Alternative Chloramphenicol (500mg)QID x7-15 days
Enteric Fever (OPD)	T. Cefixime 400 mg TDS for 14 days	Alternative T. Azithromycin (1gm)OD for 5 days
Enteric Fever(IPD)	Ceftriaxone (4gm/day)IV for 7-14 days	

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Table 10 Pediatric Infections

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2	Patient Type 3
Pneumonia AGE: 3 weeks to 3 months	Community acquired Pneumonia Ceftriaxone 100mg/kg/d od or Cefotaxime 150mg/kg/d tds x 10-14 days and *Azithromycin 10mg/kg/day x	Either Type II or Early HAP/VAP Piperacillin-tazobactam 300 mg/kg/d qid	Either Type III or late HAP/VAP, IV Meropenem (60-120 mg /kg/day divided 8 hrly) plus Vancomycin (40-60 mg/ kg/ day divided 6-8 hrly IV Meropenem (60-120 mg /kg/day divided 8 hrly) plus Vancomycin (40-60 mg/ kg/ day divided 6-8 hrly. Add Fluconazole 6-12 mg/kg/day or amphotericin B (if renal dysfunction)
	AGE: 4 months to 5 years Lobar pneumonia/effusion Ceftriaxone 100mg/kg/d od with Cloxacillin 100-200mg/kg/d Bronchopneumonia without effusion Ampicillin 200mg/kg/d qid days*consider adding macrolide (azithromycin,) to cover Pertussis in partially unimmunized with DPT	Piperacillin-tazobactam 300 mg/kg/d Qid plus Vancomycin (40-60 mg/ kg/ day divided 6-8 hrly Ceftriaxone 100mg/kg/d od Or Piperacillin-tazobactam 300 mg/kg/d qid	Same as above
Meningitis	Community acquired	Either type II/post neurosurgical meningitis	Either type II/III or post shunt infection
Age > 3 months	Cefotaxime 200 mg/kg/d qid/or Ceftriaxone 100mg/kg/d od/bd plus Vancomycin*60mg/kg/d	IV Meropenem (120 mg /kg/day divided 8 hrly) plus Vancomycin(60	IV Meropenem (120 mg /kg/day divided 8 hrly)/ plus Vancomycin 60mg/kg/d qid with or without rifampin 10 mg/kg (PO)

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Dwivedi Medical Director
--	---	---	--



	qid*Discontinue Vancomycin if rapid latex agglutination negative for S. pneumoniae, or positive for N. meningitidis, or H. influenzae	mg/ kg/ day divided 6 hrly +/- rifampin 10 mg/kg(PO) q12h	q12h x 7-10 days after shunt removal Consider additional Intraventricular therapy Vancomycin 10mg or Gentamicin 1-2 mg or Polymixin B 2mg or Colistin 10mg[1mg = 12,500 units]
Urinary Tract Infection			
Cystitis	Co-trimoxazole 8-10 mg/kg/d of trimethoprim bd OR Amoxy-clav 30-40 mg/kg/d bd OR Cefixime 8-10 mg/kg/d od		
Pyelonephritis	<p>Uncomplicated Amoxy-clav 30-40 mg/kg/d bd OR Ceftriaxone100mg/kg/d od OR Cefotaxime150mg/kg/d tds</p> <p>Complicated: Ceftriaxone100mg/kg/d od OR Cefotaxime150mg/kg/d tds OR Piperacillin- tazobactam 300 mg/kg/d tds/qid +/- Amikacin 15- Piperacillin-tazobactam 300 mg/kg/d tds/qid +/- Amikacin 15- 20mg/kg/d od X10-14 days</p>	Piperacillin-tazobactam300 mg/kg/d tds/qid Or Meropenem120mg/kg/d	<p>Same as for type II</p> <p>Same as for type II</p>

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



<p>HEENT Infections Orbital cellulitis</p>	<p>Cloxacillin 200mg/kg/d plus either Cefotaxime 150mg/kg/d tds or Ceftriaxone 100mg/kg/d od/bd x 10-14 days</p>	<p>Piperacillin-tazobactam 300 mg/kg/d tds/qid plus Vancomycin 60mg/kg/d qid</p>	<p>IV Meropenem (120 mg /kg/day divided 8 hrly)/ plus Vancomycin 60mg/kg/d qid</p>
<p>Bone and Joint Infections Acute Osteomyelitis/s eptic arthritis</p>	<p>Cloxacillin 200mg/kg/d plus either Cefotaxime 150mg/kg/d tds or Ceftriaxone 100mg/kg/d od/bd x 10-14 days</p>	<p>Vancomycin 60mg/kg/d qid or Clindamycin 20-40 mg/kg/d tds/qid Plus either Cefotaxime 150mg/kg/d tds or Ceftriaxone 100mg/kg/d od/bd</p>	<p>IV Meropenem (120 mg /kg/day divided 8 hrly)/ plus Vancomycin 60mg/kg/d qid or Clindamycin 20-40 mg/kg/d tds/qid</p>
<p>Osteochondritis</p>	<p>Piperacillin-tazobactam 300 mg/kg/d tds/qid or combination therapy with cloxacillin 200mg/kg/d plus Ceftazidime 100mg/kg/d tds 7-10 days after surgery</p>		
<p>Skin and soft tissue infections</p>	<p>Cloxacillin 200mg/kg/d or Cefazolin 60-100mg/kg/d or Clindamycin 20-40 mg/kg/d tds/qid x 7-10 days</p>	<p>Vancomycin 60mg/kg/d qid</p>	<p>Piperacillin- tazobactam 300 mg/kg/d tds/qid or IV Meropenem (120 mg /kg/day divided 8 hrly plus Vancomycin 60mg/kg/d qid</p>

<p>Prepared by Dr M Kumar Prof Microbiology</p>	<p>Checked by Dr M S Modak Prof Microbiology</p>	<p>Verified by Dr AK Verma Quality Department</p>	<p>Approved by Dr SK Talwani Medical Director</p>
---	--	---	---



<p>Animal bite wounds (dog / cat)</p>	<p>Amoxicillin/clavulanate 50mg/kg/d tdsi.v or p.o</p>	<p>Alternatives Piperacillin300mg/kg/d qid 7-10 days <u>Penicillin allergy</u> Clindamycin20-40mg/kg tds/qid plus TMP /SMX 80mg/kg/ bd X 7-10 days (dog bites); or cefuroxime 20-30mg/kg/d x 7-10 days (cat bites)</p>	<p>NA</p>
<p>Vascular catheter associated Infections</p>		<p>Piperacillin-tazobactam300 mg/kg/d tds/qid + Vancomycin60mg/kg/d qid</p>	<p>Meropenem 120mg/kg/d tds plus Vancomycin 60mg/kg/d qid</p>
<p>Severe Sepsis/septic shock</p>	<p>Cefotaxime 150 mg/kg/day divided 6-8 hrly OR Ceftriaxone 100 mg/kg/day divided 12 hrly +/- amikacin 15-20 mg/kg/d od</p>	<p>IV Piperacillin – Tazobactam 300-400 mg/kg/day divided 8 hrly + IV Vancomycin 45-60 mg/kg/day divided 6-8 hrly</p>	<p>IV Meropenem 80-120 mg/ kg/8 hrly + IV Vancomycin 45-60 mg/kg/day divided 6-8 hrly</p>

<p>Prepared by Dr M Kumar Prof Microbiology</p>	<p>Checked by Dr M S Modak Prof Microbiology</p>	<p>Verified by Dr AK Verma Quality Department</p>	<p>Approved by Dr SK Talwani Medical Director</p>
---	--	---	---



Table 11: Empiric Therapy of Neonatal Intensive Care Unit Sepsis and Meningitis

Diagnosis	Suspected organisms	Early onset	Late onset	Nosocomial	Community acquired	Duration
Sepsis	Klebsiella, Acinetobacter, E.coli, Enterococcus, Others :Serratia, Burkholderia, Pseudomonas, Proteus	Gentamicin (for haemodynamically stable) Piperacillin-Tazobactam (for haemodynamically unstable)	1 st line :Piperacillin-Tazobactam 2 nd line: Meropenem 3 rd line: Colistin	1 st line Piperacillin-Tazobactam 2 nd line: Meropenem 3 rd line: Colistin	1 st line :Cefotaxime and Amikacin 2 nd line:Piperacillin-Tazobactam 3 rd line: Meropenem 4 th line: Colistin	10days
Pneumonia	E coli, Klebsiella, Acinetobacter, Enterococcus, Staphylococcus (CONS) Others :Serratia, Burkholderia, Pseudomonas, Proteus	Gentamicin (haemodynamically stable) Piperacillin-Tazobactam (haemodynamically unstable)	1 st line :Piperacillin-Tazobactam 2 nd line: Meropenem 3 rd line: Colistin	1 st line Piperacillin-Tazobactam 2 nd line Meropenem 3 rd line Colistin	Ceftriaxone plus Azithromycin	7days
NEC			1 st line Piperacillin-Tazobactam and Amikacin 2 nd line Meropenem 3 rd line Colistin	1 st line Piperacillin-Tazobactam 2 nd line Meropenem 3 rd line Colistin	1 st line Piperacillin-Tazobactam 2 nd line Meropenem 3 rd line Colistin	7-10days

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



Meningitis	For early onset: E coli, GBS, enteric bacilli, listeria, streptococcus, H influenza, Neisseria meningitides. For late onset: Klebsiella, Acinetobacter, E.coli, Enterococcus, Staphylococcus (CONS) Others :Serratia, Burkholderia, Pseudomonas, Proteus	1 st line: Cefotaxime plus Gentamicin 2 nd line: Meropenem	Meropenem	Meropenem	Ceftriaxone /cefotaxime	Gram Positive: 14-days Gram negative: 21 days# #Ventriculitis/Brain abscess: 6-8 weeks
UTI	Enterococcus, E coli, Enterobacter		1 st line: Piperacillin-Tazobactam 2 nd line: Meropenem 3 rd line: Colistin	1 st line Piperacillin-Tazobactam 2 nd line: Meropenem 3 rd line: Colistin	Amikacin	10days
Skin and soft tissue infection	Staphylococcus		1 st line:Cloxacillin 2 nd line:Vancomycin	Vancomycin	Cloxacillin	7days
Arthritis	Staphylococcus, Klebsiella		1 st line Piperacillin-Tazobactam 2 nd line Meropenem	1 st line: Piperacillin-Tazobactam 2 nd line: Meropenem	Ceftriaxone plus Vancomycin	Culture Negative: 2weeks

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



			3 rd line Colistin	3 rd line Colistin		Culture positive: 3 weeks
Osteomyelitis	Staphylococcus, Gram Negative Bacilli		1 st line Piperacillin- Tazobactam 2 nd line Meropenem 3 rd line Colistin	1 st line Piperacillin- Tazobactam 2 nd line Meropenem 3 rd line Colistin	Ceftriaxone plus Vancomycin	4 weeks
Catheter related Infection	Staphylococcus(CON S), S.aureus, Gram negative bacteria		1 st line: Vancomycin and Amikacin 2 nd line:Piperacil lin-Tazobact 3 rd line: Meropenem 4 th line Colistin			10days
Fungal infection	Candida albicans and Candida Non albicans		Amphotericin B or Fluconazole (depending on Antifungal susceptibility report)			Depending on location

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwar Medical Director
--	---	--	---



Table : 12 Empiric therapy of Ophthalmic infections

Sr. No	Category	Suspected Organisms	First Line	Alternative
1	Bacterial conjunctivitis	S aureus and albus H Aegyptius H Influenzae, C diphtheriae	Topical Moxifloxacin 0.5% eyedrops 3-6 times per day Tobramycin eye ointment at bed time Penicillin eye drops 10,000 units/ml	
2	Bacterial Keratitis	Pseudomonas, S.aureus Pneumococcus N gonorrhoeae	Moxifloxacin eye drops 0.5% 1 hourly Fortified Tobramycin eye drops	Fortified Vancomycin eye drops Amikacin eye drops
3	Fungal Keratitis	Aspergillus, Fusarium, Candida albicans	Natamycin eye drops 6 times a day Itraconazole eye drops /ointment at bed time Tablet Fluconazole 150mg twice a day & eye drops 4-6 times per day Nystatin eye ointment	Amphotericin B eye drops Voriconazole eye drops Intracameral Amphotericin B
4	Viral Keratitis	H Simplex H Zoster	Acyclovir Tablet 800mg 5 times a day and ointment 5 times a day Gancyclovir ointment	Tablet Valacyclovir 1000mg 3 times a day
5	Endophthalmitis	S aureus Sepidermidis Streptococcus Pseudomonas H Influenzae Candida /fusarium	Intravitreal Vancomycin 1 mg /0.1 ml and Amikacin 400microgrames /.ml Intravitreal Amphotericin B	Intravitreal Vancomycin 1mg /0.1ml and Ceftriaxone 2.25mg/0.1ml
6	Orbital cellulitis	Staphylococci Mucormycosis/Aspergillosis	Intravenous Piperacillin and Tazobactam 4.5g twice a day Intravenous Metronidazole 100ml 3 times a day Intravenous Amphotericin B	Intravenous Ceftriaxone
7	Acute Dacryocystitis	Staphylococcus, Streptococcus, Pneumococcus	Tablet Amoxicillin and Clavulanic acid 625 mg twice a day Moxifloxacin eye drops 0.5% 3-6 times a day	

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--

**Table 13: ENT Infection**

Name of condition	Patient Type 1 (Community acquired)	Patient Type 2	Patient Type 3
Acute infection like acute membranous tonsillitis, ASOM, Acute epiglottitis without complication	Inj Ampicillin 1 gm q6h Amoxicillin +clavulanic acid 1.2 gm q8h	-	-
Acute infection with complications like acute mastoiditis, Quinsy	Addition Of aminoglycoside for gram negative coverage and metronidazole for anaerobic coverage	-	-
Chronic infection without complication like CSOM, chronic sinusitis	Amoxicillin +clavulanic acid 1.2 gm q8h IV Ceftriaxone 1 gm q12h IV	ID/ Medicine consult	ID/Medicine consult
Chronic infection with complications like meningitis, orbital cellulitis, brain abscess	Inj Ceftriaxone+ inj amikacin + inj metronidazole	ID/ Medicine consult	ID/ Medicine consult

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Table 14: Surgical site infection

Name	Type 1	Type 2	Type 3
Head & Neck	Ceftriaxone 1gm q12h IV + Metronidazole Or PIP-TZ 4.5 gm q6h IV If MRSA suspected Add Vancomycin 1gm IV q12h If CNS infection Ceftazidime 2 gm q8h IV instead of Ceftriaxone/PIP-TZ	Meropenem 2gm q8h IV + Vancomycin 1 gm q12h IV	If fungal infection suspected Ampho B If VRE suspected Linezolid If XDR or PDR Gram negative infection suspected Colistin 4.5MUBD If CNS infection Add intrathecal antimicrobials as above
Other infections Sternal infections Chest Abdominal Perineal	Ceftriaxone 1gm q12h IV + Metronidazole Or PIP-TZ 4.5 gm q6h IV If MRSA suspected Add Vancomycin 1gm IV q12h	Meropenem 2gm q8h IV + Vancomycin 1 gm q12h IV	If fungal infection suspected Ampho B If VRE suspected Linezolid If XDR or PDR Gram negative infection suspected Colistin 4.5MUBD If clostridium difficile colitis or sepsis suspected Oral Vancomycin 250 mg q6h + Metronidazole 500 mg q8h IV

Note:

Surgical debridement is almost always necessary. Any graft, device or foreign body must be removed.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	---	--	--



Table 15: Catheter related blood stream infections (CRBSI)

Name	Type 1	Type 2	Type 3
Peripheral catheter		Cloxacillin 1 gm q6h IV	Ceftriaxone 1gm - q12h IV
Central venous catheter (short term) Dialysis catheter (short term)	-	+ Meropenem 2gm q8h IV Vancomycin 1 gm q12h IV	Meropenem 2gm q8h IV + Vancomycin 1 gm q12h IV
Dialysis catheter (long term) Hickman or other implanted catheter (long term)			If fungal infection (Non-Albicans Candida suspected) Ampho B iv Or Caspofungin 70 mg IV q24h flowed by 50 mg If VRE suspected Linezolid If XDR or PDR Gram negative infection suspected Colistin 4.5MU BD

Note:

Change catheter if signs of thrombophlebitis are present

Catheter cultures and blood cultures to be sent as per HICC protocol.

Catheter maybe kept in situ pending culture reports especially if CRBSI not strongly suspected and no other IV access is available

Remove catheter immediately if local signs of suppuration present or if central venous catheter and blood cultures are positive

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	--	---	--



DEFINITIVE THERAPY ONCE THE CAUSATIVE ORGANISM IS IDENTIFIED

It is vital to send cultures before empiric antimicrobials are prescribed. Once cultures results are available the next steps are

1. Decide whether the organism grown is a colonizer or an actual pathogen. Evaluate carefully if the site from which culture has been sent has active infection either from clinical signs or from elevated WBC counts or radiological evidence.
2. Don't treat colonizing organisms, Consult microbiology for the decision
3. Choose the simplest antimicrobial class to which the organism shows susceptibility
4. If the cultures show intermediate susceptible or multidrug resistant organism, consult infectious disease specialist for choice of appropriate antimicrobial.
5. Linezolid should be given only in culture confirmed MRSA infections after consultation with ID physician.
6. Levofloxacin is reserved for use in culture confirmed pulmonary infections only
7. Do not continue therapy beyond indicated duration. If the duration is to be exceeded then the clinician will justify the same and endorse it in the clinical notes

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



**Table 16. Antimicrobial Prophylaxis for Surgery
Clean and Clean Contaminated cases**

Procedure	Antimicrobial
Clean surgeries (example: elective hernia repair, breast surgeries)	Cefazolin / Cefuroxime
Orthopedic surgery	Cefazolin / Cefuroxime (add amikacin and metronidazole in open fractures)
Cardiovascular / vascular surgery	Cefazolin / Cefuroxime
Neurosurgery	Cefazolin / Cefuroxime
Ophthalmic surgery	Topical quinolone. Systemic- Cefazolin / Cefuroxime
Head, neck and ENT surgery	Cefazolin / Cefuroxime/Amoxycylav (Ceftriaxone in cases involving dural exposure)
Gastroduodenal	Cefuroxime / Cefazolin
Appendicular / Colorectal surgery Biliary	Cefuroxime / Cefazolin and Metronidazole Cefuroxime / Cefazolin/ cefoperazone-sulbactam
Abdominal / Vaginal hysterectomy / Caesarian section/MTP/Sterilization/Laparoscopic procedures	Cefazolin / Cefuroxime +Metronidazole
Urologic surgery	Cefuroxime (or as guided by urine culture)
Ophthalmology Intraocular surgeries under LA Surgeries under GA (Clean Surgeries)	Topical Quinolone eye drops IV Inj. Cefazoline / Cefuroxime Topical Quinolone eye drops

Note: Preoperative dose of antimicrobial is to be given 15 to 60 minutes before incision

- Dose of Cefazolin 2 gm IV
- Dose of Cefuroxime 1.5 gm IV
- Dose is to be repeated if surgery > 4 hours
- If penicillin allergy: consider either clindamycin or Vancomycin (infusion to be started more than 60 min before incision for vancomycin in view of prolonged infusion time required) (WHO Guidelines on surgical prophylaxis 2018)
- Antimicrobial prophylaxis must not be continued for more than 24 hours after surgery
- In case patient is already on antimicrobials the same antimicrobial should be adjusted for the period of surgery.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--

**Table 17 : Pediatric surgery**

Procedure	Antimicrobial	Duration
Urosurgery	Amoxiclav	3 days
Genitourinary: Clitoroplasty Hypospadias, vaginoplasty (plain) Vaginoplasty (use of bowel)	Amoxiclav Ceftriaxone + Metronidazole	5days 1 day prior
Hirschprung ARM pull-through	Ceftriaxone + metronidazole	3 days
Esophageal procedures	Amox-clav	
Biliary tract	Cefoperazone -sulbactam	3 days
Appendix : Nonperforated Perforated	Cefuroxime Ceftraxone + Metronidazole Piptaz if required	1 dose
Paediatric Tumours	Cefuroxime	3 doses

Clean surgeries:

Procedure	Antimicrobial	Duration
Circumcision, Orchiopexy	Cefuroxime	Single dose
Herniotomy, LN biopsy, Thyroglossal cyst Lipoma, small lumps	No antimicrobials	-

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwar Medical Director
--	---	--	---



Table 18 : Antimicrobial prophylaxis for GI endoscopy

Antimicrobial prophylaxis and/or treatment to prevent local infections					
	Patient condition	Procedure contemplated	Goal of prophylaxis	Antimicrobial	Periprocedural antimicrobial prophylaxis
	Bile duct obstruction in absence of cholangitis	ERCP with complete drainage	Prevention of cholangitis	Nil	Not recommended
	Bile duct obstruction in absence of cholangitis	ERCP with incomplete drainage	Prevention of cholangitis	Ceftriaxone	Recommended; continue antimicrobials after procedure X 3 days
	Solid lesion in upper GI tract	EUS-FNA	Prevention of local infection	-	Not recommended
	Solid lesion in lower GI tract	EUS-FNA	Prevention of local infection	-	Not recommended
	Pancreatic cysts	EUS-FNA	Prevention of cyst infection	Ceftriaxone+ Metronidazole	Three doses
	All patients	Percutaneous endoscopic feeding tube placement	Prevention of peristomal infection	Cefazolin/Cefuroxime	Recommended single dose
	Cirrhosis with acute GI bleeding	Required for all patients regardless of endoscopic procedures	Prevention of infectious adverse events and reduction of mortality	Ceftriaxone	On admission
	Synthetic vascular graft and other nonvalvular cardiovascular devices	Any endoscopic procedure	Prevention of graft and device infection	-	Not recommended
	Prosthetic joints	Any endoscopic procedure	Prevention of septic arthritis	-	Not recommended
	Peritoneal dialysis	Lower GI endoscopy	Prevention of peritonitis	Ceftriaxone + metronidazole	Suggested
EUS-FNA, EUS-guided FNA.					

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Table -19 FEBRILE NEUTROPENIA

Febrile Neutropenia-definition

- Neutropenia-ANC<500/mm³and expected to fall below 500/mm³ in 48hrs
- Fever-single oral temperature of 38.3oC(101oF) on one occasion or 38oC (100.4oF) on atleast 2 occasions (1 hour apart)
- Neutropenic patients may not have usual signs of infection. Redness, tenderness and fever may be the only signs.

Protocol:

- Critical examination of areas usually harboring infections, including but not limited to, oral cavity, axillary region, scalp, groin, perineal region.
- Send blood Cultures 2 sets (each bottle 10ml x 4 bottles)
- Other relevant investigations: urea, creatinine, ALT, urine culture ,Chest Xray, separate culture from central line, etc.

Patient-Haemodynamically stable

- Blood culture 2 sets
- Start IV Cefoperazone sulbactam 1.2gm IV 8 hourly
- No need to add glycopeptides in the initial regimen (except in specific situations, given below)

Patient-Haemodynamically unstable

- Start BL-BLI agent(Cefoperazone-Sulbactam 1.2gm IV 8 hourly/ piperacillin- tazobactam 4.5gm IV 8 hourly) OR Carbapenem (meropenem 1gm IV 8 hourly/imipenem 500mg IV 6 hourly/doripenem 500mg IV 6 hourly)
- No need to add glycopeptides in the initial regimen (except in specific situations, givenbelow)

Reassess after 48 hours:

If blood cultures are negative, haemodynamically stable but still febrile

- Reculture blood
- Add amikacin 500mg IV BD for 3days
- Add colistin (instead of amikacin) if indicated (see below)

If blood cultures are negative, haemodynamically unstable but still febrile

- Inj Colistin (+/-Carbapenem) + glycopeptides + Echinocandin/ L-AmphoB

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Blood culture growing Gram negative bacilli

- Patient afebrile- continue the empirical antibiotic till antibiotic sensitivity is available.
- Rationalise as per susceptibility profiles

When to add glycopeptides?

1. Haemodynamic instability, or other evidence of severe sepsis, septic shock or pneumonia
2. Colonisation with MRSA or penicillin-resistant S. pneumonia
3. Suspicion of serious catheter-related infection e.g. chills or rigors within fusion through catheter and cellulitis around the catheter exit site
4. Skin or soft-tissue infection at any site
5. Positive blood culture for gram-positive bacteria, before final identification and susceptibility testing is available
6. Severe mucositis

When to add empirical colistin in febrile neutropenic patients?

1. Haemodynamic instability.
2. Colonisation with carbapenem resistant gram-negative bacteria.
3. Previous infection with carbapenem resistant gram-negative bacteria.
4. GNB in blood, sensitivity pending, persistent fever with haemodynamic instability.

Empirical Antifungal Therapy

- No response to broad spectrum antibiotics (3-5days)- add L-AmphoB/echinocandin
- When a patient is located at a remote area and may not have access to emergency healthcare services, febrile neutropenia can be life threatening. Under such circumstances, availability of broad-spectrum oral antibiotics with the patient can help them gain time to reach emergency healthcare service.

Useful tips

- Febrile after 72hrs- CT chest and consider empirical antifungal.
- If fever persists on empirical antibiotics, send two sets blood cultures/day for 2 days
- Send further cultures if clinical deterioration
- Unexplained persistent fever in otherwise stable patient doesn't require change in empirical antibiotic regimen.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Continue the regimen till ANC is >500cells/mm³

- If glycopeptides started as a part of empirical regimen, STOP after 48hrs, if no evidence of Gram positive infection
- Antibiotic treatment should be given for at least seven days with an apparently effective antibiotic, with at least four days without fever.
- Once Neutrophil count has recovered, with no culture positivity and hemodynamically stable; antibiotics can be stopped and patient observed, even if remains febrile. Evaluate for fungal infection, if at risk.

Antiviral prophylaxis

- For HSV IgG positive patients undergoing allo-HSCT or leukemia induction needs acyclovir prophylaxis
- All patients being treated for cancer need to receive annual influenza vaccination with an inactivated vaccine.
- Neutropenic patients presenting with influenza like illness should receive empirical treatment with neuraminidase inhibitor.

Antifungal prophylaxis

- a) Induction chemotherapy of Acute Leukemia: Posoconazole
- b) Post allo BMT

Pre engraftment:

Voriconazole/ echinocandin

Post engraftment:

Posoconazole

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	---	--	--



Table - 20 Cardiac Device implant Prophylaxis

A risk assessment score is proposed with one point for each

a) Diabetes mellitus	b) Heart failure
c) Oral anticoagulation therapy	d) Chronic corticosteroid use
e) Renal insufficiency/failure	f) Prior CIED infection
g) Presence of more than two leads implanted	h) Presence of epicardial leads
k) Use of temporary pacemaker	l) Replacement, or upgrade procedure.

According to the score, patients will be stratified in two groups and the **same documented in the case notes:**

(a) low infective risk (score <3)
 (b) high infective risk (score ≥3)

Patients in the 'low-risk' group will treated with only two doses of antibiotics, both intravenous, of whom the first 15-60min before skin incision and the second after 8 hours. Patients in the 'high-risk' group will get intravenous prophylaxis for two full days (of whom the first administration 15-60min before skin incision

Antibiotic prophylaxis will be amoxicillin/clavulanic acid unless the patient had a history of allergic reactions to penicillin. The dosage will be dependent on renal function:

(a) intravenous amoxicillin/clavulanic acid 2/0.2 g in patients with creatinine clearance (CrCl) >30 mL/min and 1/0.2 g in patients with CrCl <30mL/ min
 (b) oral amoxicillin/clavulanic acid 875/125 mg every 8 h in patients with CrCl >30 mL/min, and 875/125 mg every 12 h in patients with CrCl <30mL/min.
 (c) In case of penicillin allergy Intravenous clindamycin day for 2 doses for the low risk group and two days for the high risk group.


Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Appendix 1

Commonly used antimicrobials

CLASSIFICATION OF ANTIBIOTICS



Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwar Medical Director
--	---	--	---



Table 21 : Spectrum of commonly used antimicrobials:

Antimicrobial Class	Name	Organisms	Indication & Dose	Side effects
Penicillins				Allergy
β-lactamase susceptible	Penicillin G Penicillin V Ampicillin. Amoxicillin (PO)	Gram +ve Gram +ve Gram +ve & Gram -ve Gram +ve	Not easily available 1-2 gms q6h 500 mg q8h	
β – lactamase resistant	Cloxacillin	Gram +ve	0.5-1gm q6h	
β-lactam/ β-lactam inhibitor combination	Piperacillin-tazobactam. Ampicillin-sulbactam. Amoxicillin-clavulanate (IV)	ESBL Gram -ve organisms ESBL Gram -ve organisms Gram +ve & Haemophilus. influenzae	4.5 gm q6h as infusion 1 gm q6h 1.2 gm q8h	
Cephalosporins				
1 st Generation	Cefazolin (IV) Cephalexin (PO)	Gram +ve	1gm q8h 500 mg q8h	
2 nd Generation	Cefadroxil (PO) Cefuroxime (PO & IV)	Gram +ve Gram +ve	500 mg q12h 750 mg q8h	
3 rd Generation	Cefotaxime Ceftriaxone Ceftizoxime Ceftazidime Cefixime (PO) Cefpodoxime (PO) Cefdinir (PO)	Gram +ve & Gram -ve Gram +ve & Gram -ve Gram +ve & Gram -ve Gram +ve & Gram -ve Anti-pseudomonas	1 gram q6h 1-2gm q12h 1 gm q12h 1-2 gm q8h 200 mg q12h	

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Antimicrobial Class	Name	Organisms	Indication & Dose	Side effects
Cephalosporin Plus beta lactamase inhibitor	Cefoperazone /sulbactam	Anti-pseudomonas	q12h 1.5 gm – 3gm q12h	
Aminoglycosides	Streptomycin Kanamycin Gentamicin Amikacin Tobramycin Netilmicin	Gram –ve Gram –ve Gram –ve Gram –ve Gram –ve Gram -ve	0.75 -1gm q24h 3mg/kg q24h 13mg/kg q24h 3mg/kg q24h 5mg/kg q24h	Deafness Vertigo Muscle weakness
Quinolones	Nalidixic acid Norfloxacin Ciprofloxacin Ofloxacin Levofloxacin Moxifloxacin		1 gm q6h 400 mg q12h 500 mg q12h 200 mg q12h 750 mg q24h 400 mg q24h	Seizures
Extended spectrum Carbapenems Imipenem-cilastatin Meropenem Doripenem Ertapenem		Gram +ve except MRSA, ESBL Gram –ve except Stenotrophomonas, Burkholderia, Corynebacterium, Enterococcus faecium not covered Does not cover Pseudomonas, Acinetobacter & Enterococcus	0.5gm -1gm q6h 1 – 2 gm q8h 1gm q24h	Seizures
Polymyxins Polymyxin B Colistin		ESBL, Metalloproteinase producing Gram –ve	Colistin 4.5MUBD (loading dose of 9MU is mandatory)	Muscle weakness Renal toxicity
Lincosamide Clindamycin		Gram +ve and anaerobes	600mg q8h	C. difficile colitis

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Antimicrobial Class	Name	Organisms	Indication & Dose	Side effects
Glycopeptides Vancomycin Teicoplanin		MRSA	1 gm q12h 400 mg q24h	Renal toxicity
Oxazolidinedione Linezolid		VRE	600 mg q12h	Thrombocytopenia
Lipopeptides Daptomycin		MRSA	4-6mg/kg q24h	
Antifungals Fluconazole Voriconazole Caspofungin Anidulafungin AmphoB aqueous AmphoB colloidal AmphoB liposomal		Candida albicans Aspergillus Non albicans candida Non albicans candida Broad spectrum covers all above + Mucor etc	400 mg q12h 6mg/kg q12h first day then 4mg/kg 70mg IV then 50 mg q24h Refer product insert	

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwani Medical Director
--	---	--	--



Appendix 2

Duration of therapy for various clinical conditions

Short course therapy is equally effective			
Condition	Short Course (Days)	Long Course (Days)	Outcome
Chronic Bronchitis and obstructive pulmonary disease, acute exacerbations*	<=5	>=7	Equivalent Most patients do not require antimicrobials at all
Intra-abdominal infection	4	10	Equivalent
Neutropenic fever	Until afebrile and stable	Until non neutropenic	Equivalent
Osteomyelitis, Chronic	42	84	Equivalent
Pneumonia community acquired	<=8	10-15	Equivalent
Pyelonephritis	5-7	10-14	Equivalent
Skin Infections (Cellulitis, Major abscesses, wound infections)	5-6	10-14	Equivalent
Sinusitis, acute bacterial	5	10	Equivalent

Reference : Harrison's principle of Internal Medicine 20th ed

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



Appendix 3 Antimicrobial Agent Form

	BHARATI VIDYAPEETH UNIVERSITY MEDICAL COLLEGE HOSPITAL AND RESEARCH CENTRE
	ANTIMICROBIAL AGENT FORM
	BHRC/HIC/F23(V-2)

List of high end antibiotics:
 1. Carbapenems 2. Piptaz 3. Levofloxacin 4. Colistin, 5. Polymyxin B , 6. Fosfomycin/daptomycin 7. Teicoplanins 8. Vancomycin,
 9. Tigecycline/ Minocycline 10. Teicoplanins 11. Linezolid 12. Echinocandins 13. Voriconazole/ Posaconazole 14. Amphotericin B

Ward : ICU I/II/III (Surg/ Ortho) Med 3/6/10/11 Other Wards: _____

Bed No : _____ Clinician/Unit Head : _____

Date of Admission: [][][][][][] Date of filling: [][][][][][]

Paste Patient Barcode here

Confirmed Diagnosis: _____ Patient type : 1 2 3

Suspected cause/ site of infection: _____ Wt of patient _____ kg.

Date of Surgery: [][][][][][]

Time of incision [][][][] Time of 1st dose [][][][] Time of 2nd dose [][][][] Duration of surgery [][][][] HRS

AMA Details: 1. High End Antibiotics 2. Other Antibiotics

AMA(Generic name)	E/D/SP*	Route , Freq and Dose (mg/g)	Start Date	Day 1	Day3	Day7	IV to Oral	Stop Date	CHANGE OF AMA (remarks)
				(TEMP, TLC ,PCT, SERUM CREATININE)					

*=E- Empirical; D- Definitive; SP- Surgical Prophylaxis

Device in situ : 1. Central Line 2. Peripheral line 3. HD Catheter 4. Urinary Catheter 5. ETT/TT 6. Drains

Date of insertion _____ Date of removal _____

Culture/ Sensitivity Investigations done: YES / NO; If YES; Date of Report:

Specimen taken (B/P/U/Resp/CSF)* and Date(DD/MM/YY) (1BEFORE,2AFTER)	Microorganism isolated Date received(DD/MM/YYYY)	Sensitivity Pattern (Name of imp antibiotics)	Change of AMA Y/N	Response after change Y/N

*=B- Blood ; P- Pus ; U- Urine; Resp – Sputum,ETT,BAI ; CSF- Cerebro spinal fluid

Collected Data by : _____

Prescriber sign : _____
(in case of High end Antibiotic)

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Dwivedi Medical Director
--	--	---	---



Appendix 4

Operative Procedure for Sample Collection

1. URINE CULTURE

(a) Items required

The following items should be available as preparation for collecting a catheter specimen of urine for analysis:

- Sterile gloves;
- Alcohol-saturated swab;
- Gate clip or non-traumatic clamps;
- Sterile Universal specimen container;



(b) Collection of urine sample from indwelling catheter:

Use sterile precautions

Clean hands using hand rub

Clamp the distal portion of the catheter.

Disconnect urine bag.

Disinfect distal portion of the catheter using 70% alcohol swab Allow it to dry thoroughly.

Do not allow the distal end of the catheter to touch body or clothes of the patient.

After 10 minutes release the clamp and collect urine in a sterile urine collection container



(c) Urine must be transported to the lab without delay. If delay is unavoidable, sample should be stored in the refrigerator at 4° C.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department	Approved by Dr SK Talwar Medical Director
--	---	--	---



2. BLOOD CULTURE

(a) Hands should be clean and dry and gloved before taking sample.




(b) Prepare Blood culture vials by proper labeling

(c) The culture bottle should be brought to room temperature before the sample inoculation and should not be refrigerated after inoculation.

(d) Remove flip-off caps and wipe the rubber bung with a 70% alcohol swab

(e) Disinfect skin at venepuncture site by wiping with 70% alcohol in a circular motion from centre to periphery and then with 1% iodine. Wait for at least one minute, allow to dry before venipuncture.



(f) After venepuncture carefully withdraw the needle and compress the venepuncture site.

(g) The number of sets to be collected is as follows.

(i) Febrile episode: 2 bottles from separate sites within 10 min

(ii) Acute endocarditis: 6 bottles from separate sites at 30 min intervals

(iii) Central Line related : 1 bottle drawn from the central line and 1 bottle from a peripheral venipuncture site

(h) The various culture bottles for Automated Blood Culture System and the volume of blood to be added to them are as under.

(i) BacT/Alert/BACTEC Aerobic (30 ml): 10 ml blood (optimal)


(ii) BacT/Alert/BACTEC Paediatric (20 ml): 4 ml blood (optimal)

Prepared by	Checked by	Verified by	Approved by
Dr M Kumar Prof Microbiology	Dr M S Modak Prof Microbiology	Dr AK Verma Quality Department of Med Dir	Dr SK Talwani Medical Director



3. ENDOTRACHEAL/TRACHEOSTOMY ASPIRATE CULTURES

- (a) The collection of endotracheal aspirate will be performed by the Anaesthetist or the Intensivist.
- (b) Do not use a swab for collecting sample.
- (c) Use a BAL collection trap to collect the endotracheal aspirate.
- (d) The aspirate is to be drawn as such in case it is less tenacious or alternatively, 5 ml of sterile saline is instilled and withdrawn immediately through the disposable sterile endotracheal aspiration catheter.
- (e) The material is to be sent to laboratory within the next one hour.



4. PUS CULTURE

- (a) Clean surface of wound or abscess with 70% alcohol and allow to dry; aspirate pus or fluid from the depth of the wound in a disposable syringe, place in a sterile container and send to the laboratory.



- (b) Cotton swab to be used only if pus cannot be collected in sterile bottle or syringe.



5. Body fluids

Sample-Amniotic fluid, Synovial fluid, pericardial fluid, Pleural fluid, peritoneal fluid

Sterile fluids are usually collected by a trained, qualified physician.

Aseptically collect at least 1 ml of fluid into a new disposable sterile container.

6. Bronchoalveolar Lavage

BAL should be collected under aseptic conditions preferably with a protected specimen brush if available. The material should be collected in a BAL trap container.

7. High vaginal swabs (HVS)

Use speculum to separate the vaginal walls.

Wipe away any excess cervical mucus with a sterile cotton swab.

Use the sterile swab stick to sample as high as possible in the vaginal vault.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--



References:

1. National treatment guidelines for antimicrobial use in infectious disease, version 1 (2016)
2. ICMR Treatment guidelines for antimicrobial use in common syndromes, 2022 2nd edition
3. ICMR Antimicrobial Stewardship Program Guideline - 2018
4. Antimicrobial stewardship programmes in health-care facilities in low-and middle-income countries: a WHO practical toolkit 2019
5. Antimicrobial stewardship: Systems and processes for effective antimicrobial medicine use. NICE guideline 2015 (updated Jan 2018)
6. IDSA : New Antibiotic Stewardship Guidelines Focus on Practical Advice for Implementation 2016
7. Harrison’s Principles of Medicine 20th ed
8. ASGE guidelines for antimicrobial prophylaxis in GI Endoscopic procedures
9. Urinary Tract Infections in Pregnant Individuals. Obstet Gynecol. 2023 Aug 1;142(2):435-445. doi: 10.1097/AOG.0000000000005269. PMID: 37473414.
10. Blomstrom-Lundqvist C, Traykov V, Erba PA, Burri H, Nielsen JC, Bongiorno MG et al. European Heart Rhythm Association (EHRA) international consensus document on how to prevent, diagnose, and treat cardiac implantable electronic device infections-endorsed by the Heart Rhythm Society (HRS), the Asia Pacific Heart Rhythm Society (APHRS), the Latin American Heart Rhythm Society (LAHRS), International Society for Cardiovascular Infectious Diseases (ISCVID), and the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). Europace 2020;22:515–49.

Prepared by Dr M Kumar Prof Microbiology	Checked by Dr M S Modak Prof Microbiology	Verified by Dr AK Verma Quality Department of Med Dir	Approved by Dr SK Talwani Medical Director
--	---	---	--